

Antigen cross-presentation: extending recent laboratory findings to therapeutic intervention

T. W. H. Flinsenberg, E. B. Compeer,
J. J. Boelens and M. Boes

*Department of Pediatric Immunology, University
Medical Center Utrecht/ Wilhelmina Children's
Hospital, Utrecht, the Netherlands*

Accepted for publication 29 March 2011

Correspondence: M. Boes, Department of
Pediatric Immunology, University Medical
Centre Utrecht/Wilhelmina Children's Hospital,
Lundlaan 6, Post box 85090, KC.01-069-0, 3584
EA Utrecht, the Netherlands.
E-mail: m.l.boes@umcutrecht.nl

Summary

The initiation of adaptive immune responses requires antigen presentation to lymphocytes. In particular, dendritic cells (DCs) are equipped with specialized machinery that promote effective display of peptide/major histocompatibility complexes (MHC), rendering them the most potent stimulators of naive T lymphocytes. Antigen cross-presentation to CD8⁺ T cells is an important mechanism for the development of specific cytotoxic T lymphocyte (CTL) responses against tumours and viruses that do not infect antigen-presenting cells. Here, we review recent findings concerning antigen cross-presentation to CD8⁺ T lymphocytes. Specific subtypes of DCs in the mouse have been defined as being especially endowed for antigen cross-presentation, and a human homologue of these DCs has recently been described. DC vaccination strategies for the prevention and treatment of human diseases have been under investigation in recent years, but have not generally reached satisfying results. We here provide an overview of new findings in antigen cross-presentation research and how they can be used for development of the next generation of human DC vaccines.

Keywords: antigen cross-presentation, CD8 T cell, dendritic cell, HLA, human, MHC, mouse

Antigen cross-presentation to CD8⁺ T cells, a historical perspective

Dendritic cells (DCs) are key players in initiation and control of adaptive immune responses due to their exquisite ability to present antigenic fragments in the form of peptide/major histocompatibility complexes (MHC) to T cells [1–3]. Endocytosed antigens acquired from the outside environment are generally presented as peptide/class II MHC complexes, while antigens acquired from within the cell are predominantly presented as peptide/class I MHC complexes. This dichotomy raises one complication: how are DCs able to present viral or tumour peptides on class I MHC if these peptides are not endogenously produced? In 1976 a third mechanism was identified, whereby exogenous class I MHC-restricted antigens are captured by DCs, resulting in the induction of CD8⁺ cytotoxic T lymphocyte (CTL) responses [4]. This process was coined 'antigen cross-priming'. Cross-priming is important in anti-viral and anti-tumour immunity [5,6]. Mouse experiments in which non-haematopoietic cells were virally infected showed a requirement for cross-presentation by haematopoietic cells to elicit virus-specific CTL responses [5,7]. Secondly, antigen cross-presentation is

relevant to the induction of central immune tolerance in the thymus [8] and peripheral tolerance in the draining lymph node [9], a process referred to as 'cross-tolerance' [5,8]. In the 1990s, the term 'cross-presentation' was introduced to describe the antigen-presentation process underlying cross-priming and cross-tolerance. Experimental support again came from mouse model-based experiments, such as using the receptor interacting protein-membrane-bound ovalbumin (RIP-mOVA) mice that express a membrane-bound form of ovalbumin on restricted tissues including pancreatic β cells [9]. Analysis of RIP-mOVA thymus and control thymus grafted mice after being injected with OVA-specific CD8⁺ T cells from OT-I transgenic mice suggested that OVA-specific CD8⁺ T cells were lost and probably deleted after entry in the peripheral tissues [9]. Also, the inability of DCs to cross-present results in the accumulation of fully functional self-reactive CD8⁺ T cells that can cause autoimmune disease [10].

DCs, B cells, monocytes and macrophages are classified as prototypic professional antigen-presenting cells (APC) by virtue of their constitutive expression of class II MHC molecules. Professional APCs are critically important for induction of protective CD8⁺ T cell responses against normal

'self'-antigens [11], tumour antigens [11,12] and viruses [13]. As was already shown elegantly in 1996, the injection of OVA peptide-specific naive CD8⁺ T cells into non-irradiated RIP-mOVA mice results in selective presence of these T cells in the draining lymph nodes of OVA-expressing tissues (i.e. pancreas and kidney) and not other lymph nodes [11]. These data supported the notion that cross-presentation is a constitutive mechanism, whereby T cells can be primed to antigens that are present in non-lymphoid tissues that are normally not patrolled by circulating naive T cells.

The activation of CTL upon recognition of infection- or tumour-associated peptides encompasses risk to autoimmune T cell reactivity and is therefore under tight control. Under homeostatic as well as inflammatory conditions, tissue-specific DCs and, to a lesser degree, macrophages, execute peripheral tolerance control by their ability to discriminate between cross-presentation and cross-tolerization [14,15]. Also liver sinusoidal endothelial cells are capable of cross-presenting soluble exogenous antigen to CD8⁺ T cells leading to tolerance [16]. Other cell types are not yet described to have the ability to induce cross-tolerance under those non-inflammatory conditions. During infection, however, more cell types were recently identified as being able to cross-prime foreign peptides and elicit CTL responses. Thus far, B cells [17,18], neutrophils [19,20], basophils [21], mast cells [22] and endothelial cells [23] were also demonstrated to be capable of cross-presentation *in vitro*. Cross-presentation by basophils was even shown to be relevant in an *in vivo* experimental autoimmune encephalitis model [22]. However, the involvement of the other cell types in cross-presentation *in vivo* has not yet been shown, and particularly DCs appear pivotal for antigen cross-presentation in various circumstances as, for example, demonstrated by a lack of CTL responses against cell-associated antigens after depletion of DCs *in vivo* [24]. The efficiency of DCs to cross-present exogenous antigens as peptide/class I MHC *in vivo* was emphasized in a direct comparison study, where cross-presentation showed near equal efficiency as presentation of peptide/class II MHC derived from the same antigen [25].

Specific DC subsets are associated with antigen cross-presentation, and initial descriptions for these subsets are now reported in humans. Various mechanisms that facilitate cross-presentation by DC subsets were especially investigated in the last decade, mainly in mouse-based experiments. Human DC research that involves antigen cross-presentation is lagging behind. This review focuses on the mechanisms and cells that are known to be relevant for induction of effective CD8⁺ T cell responses to endocytosed antigens.

Mechanisms in DCs that facilitate antigen cross-presentation

The ability of DCs to cross-present antigen to T lymphocytes is not represented uniformly in all DC subsets. Some DC

types are more specialized in antigen transport from peripheral tissues to secondary lymphoid tissues, whereas others are non-migratory and are specialized at generation and display of peptide/MHC complexes to naive T cells that reside within lymph nodes. The role of the different subsets of DCs in antigen cross-presentation has been studied extensively in mice. DCs are characterized in the literature as lineage-marker-negative (CD3, 14, 15, 19, 20 and 56) and high expression of MHC class II molecules. Mouse DCs are further marked by expression of the integrin CD11c, and additional delineation can be made using additional cell surface markers [3,26–28].

Although some aspects of the human and mouse DC systems appear to be well conserved, other functions do not relate. In mice, a subset of resident DCs, characterized by high surface expression of CD8 α [29], is associated with the ability to cross-present exogenous (such as necrotic) antigens to CD8⁺ T lymphocytes [30–36]. The transcription factor Batf3 is crucial for the development of these CD8 α ⁺ DCs and absence of Batf3 in gene-targeted mice results in defective cross-presentation [37]. In 2010, the human equivalent of the mouse CD8 α ⁺ DCs was described. This human DC subset, characterized by the expression of BDCA-3 (CD141) [28], Clec9A [38,39] and the chemokine receptor XCR1 [40] was present in human peripheral blood, tonsils, spleen and bone marrow and represents a major human DC subset expressing Toll-like receptor-3 (TLR-3) [27,41]. Results indicate a dominant role for CD141⁺ DCs in cross-presentation of necrotic cell-derived antigens to CD8⁺ T lymphocytes [27], as well as superior cross-presentation of soluble or cell-associated antigen to CD8⁺ T cells when compared directly with CD1c⁺ DCs, CD16⁺ DCs and plasmacytoid DCs cultured from blood extracted from the same donors [40]. The role of this DC subset can now be scrutinized in experimental setups in laboratories across the globe. Although culturing from haematopoietic precursors is possible, the low frequency of naturally occurring CD141⁺ DCs [1 in 10⁴ peripheral blood mononuclear cells (PBMCs)] provides a further challenge before the ultimate goal of translation to clinical application using DCs to alter immune responses can be achieved.

Mechanisms that promote antigen cross-presentation that are inherent to immature DCs include their ability to actively control alkalization of their phagosomes [42], their low lysosomal proteolysis [43] and expression of protease inhibitors [44], thereby increasing the propensity that exogenous antigens engulfed in the phagosome lumen are cross-presented to CD8⁺ T cells [43]. However, there are also mechanisms restricted to DC subsets or to DC maturation stages, resulting in variability in cross-presentation efficiency. In some instances, cross-presentation ability by DCs correlates with expression of specific uptake receptors or proteins [45,46]. In addition, the nature of the antigen itself also creates a bias towards presentation via class I or class II MHC molecules [45].




Fig. 1. Proposed pathways of antigen cross-presentation. Yellow area (left side) relates to mechanisms described only in mice so far, whereas the gradient towards the red area (right side) depicts the transient increase in knowledge of antigen cross-presentation mechanisms in human cells. By receptor-mediated endocytosis, the antigen (red rod shape) is engulfed into a phagosome and subsequently processed in a cytosolic proteasome- (a,b,c) or endosomal protease-dependent (d) manner. For proteasome-mediated degradation the antigen is transported across the endosomal membrane into the cytosol by Sec61, accessed by the proteasome via endoplasmic reticulum (ER)–endosome fusion (c) or a delivery vesicle derived from the ER (a,b). After processing by the proteasome, possibly assisted by cytosolic peptidases, peptides either re-enter the endosomal compartment via transporter associated with antigen processing (TAP) where loading on class I major histocompatibility complex (MHC) may occur (a), or the canonical class I MHC presentation pathway in the ER (b,c). After proteolytic processing by endosomal pathway-resident proteases, peptides are loaded onto class I MHC molecules by replacing either exogenous peptide (recycling) or endogenous peptide (classical) loaded on class I MHC complexes (d). : antigen; : exogenous peptide; : endogenous peptide; : Sec61; : TAP; : MHC class I molecules; : proteasome; : proteases.

Once exogenous antigen is internalized by DCs, distinct mechanisms take place by which antigen-derived peptides are cleaved from larger antigen fragments and loaded onto the class I MHC molecules. To allow for display of exogenously acquired antigen in the form of peptide/class I MHC complexes, the antigen undergoes proteolytic processing to create an appropriate-sized fragment. Further restriction to the formation of peptide/class I MHC complexes involves the amino and carboxyl ends of the peptide to harbour charged anchor residues that complement those of the peptide-binding groove of the class I MHC molecule. Because the proteasome is demonstrated to be the main source of peptides in the classical MHC class I pathway, it is not unexpected that proteasome activity is thought to be essential for cross-presentation [13,45,47,48]. However, other reports have shown proteasome-independent processing of the exogenous protein via specific proteases [49,50]. This controversy has led to two different models, the dominant cytosolic pathway and the vacuolar pathway (Fig. 1).

The cytosolic pathway proposes that antigen is transported into the cytosol after internalization where proteasome degradation ensues, prior to transportation to the location of peptide assembly into peptide/class I MHC

molecules. Based on the mechanism used by DCs corroborated by the size-restriction of the antigen, internalization of antigens occurs by receptor-mediated endocytosis, pinocytosis (components < estimated 0.5 μm) or phagocytosis (components > estimated 0.5 μm). Upon internalization, antigens are located initially in phagosomes. These phagosomes fuse with early endosomes (characterized by a near-neutral, slightly acidic pH) and later with late endosomes (pH approximately 5.5). Accordingly, ultimate degradation into single amino acids takes place after fusion with acidic lysosomes, a route that is more prevalent in macrophages than in DCs [51]. Degradation within lysosomes occurs by proteases and hydrolases that have their enzymatic optimum close to the acidic pH found in lysosomes (pH 4–8) for antigen degradation [52], as well as degradation of cellular constituents as part of the normal cell homeostasis. The changes that occur to phagosomes in the endocytosis pathway is termed ‘phagosome maturation’. Phagosome maturation is important in regulation within the immune system, in the decision process as to whether an immune response is triggered or tolerance is established. The importance of this route is exemplified by changes in cell degradation that have been shown to result in autoimmune disease

[52,53]. For example, in a DNase II^{-/-} interferon (IFN)-IR^{-/-} mouse model, where macrophages were unable to degrade mammalian DNA and started the production of tumour necrosis factor (TNF), activation of synovial cells was observed resulting in chronic polyarthritis symptoms [53].

It can be deduced that the process of protein degradation following phagosome maturation must be tightly regulated. In DCs, the pH in phagosomes is kept near neutral (pH 7.5) for the first few hours after phagocytosis [54]. This is in stark contrast to rapid acidification that is seen in neutrophils and macrophages, where the pH drops to 5 within 30 min after phagocytosis [55]. Acidification of the phagosome, thereby increasing the lysosomal protease activity, has been shown to counteract cross-presentation in mice in a nicotinamide adenine dinucleotide phosphate (NADPH) oxidase complex 2 (NOX2)-dependent manner [54]. This indicates that DCs have a unique ability to regulate proteolytic activity in phagosomes, therefore controlling the amount of peptide destined for cross-presentation. Rab27a-dependent inhibitory lysosome-related organelles are involved in this pathway. These organelles are recruited continuously to phagosomes and limit acidification and degradation of ingested particles in DCs, thus promoting antigen cross-presentation [56,57]. Being able to interfere with phagosome acidification, thereby executing control of the rate of antigen cross-presentation, could provide new opportunities in increasing the efficacy of CTL targeting in DC vaccination.

The insulin-regulated aminopeptidase (IRAP) was implicated in antigen cross-presentation in peptide cleavage for generation of peptide substrates for class I MHC molecules [58]. IRAP was found in the early endosome of human monocyte-derived DCs and murine bone marrow-derived DCs, where it co-localized with MHC class I molecules and the mannose receptor (MR), but not with endoplasmic reticulum aminopeptidases (ERAPs). IRAP-deficient mice were capable of phagocytosis of antigen as well as presenting endogenously produced peptides, but cross-presented exogenous antigens with 50–70% decreased efficiency compared to wild-type mice [58]. IRAP-dependent cross-presentation requires active proteasome and function of the adenosine triphosphate (ATP)-binding cassette transporter family member TAP (transporter associated with antigen processing), but not lysosomal proteases. Therefore, this route of antigen cross-presentation involves cytosolic antigen degradation that is followed by peptide transport via TAP into IRAP⁺ endosomes. However, IRAP as well as MR appear dispensable for cross-presentation in murine splenic CD8 α ⁺ DCs but not mouse monocyte-derived DCs induced by inflammation, suggesting a role for these two molecules in inflammatory DCs, but not in steady-state CD8 α ⁺ DCs [59].

To allow for generation of peptides by the proteasome and cytosolic peptidases, antigen must traverse from the phagosome into the cytosol. Recent reports demonstrate that peptide transfer across the phagosomal membrane occurs via a selective, size-specific, reduction, unfolding (partial

proteolysis) and Sec61-dependent process [60–62]. Conversely, TAP transporters appear essential for peptide transport from the cytosol into a class I MHC loading compartment, as TAP knock-out mice are not capable of cross-presenting exogenously acquired viral peptides [7]. Moreover, efficient cross-presentation required TLR-4- and signalling molecule myeloid differentiation factor 88 (MyD88)-dependent relocation of TAP [63], essential for peptide loading of class I MHC, to early endosomes/phagosomes [64]. After processing in the cytosol, the generated peptides are transported via TAP either into the endoplasmic reticulum (ER), thereby entering the canonical class I MHC presentation pathway [13,47], or back into the phagosomal pathway [64–66]. The latter situation is likely to contribute to a rapid cross-presentation, as all necessary components are in a separate class I MHC loading-competent compartment that is distinct from the ER. Peptides generated locally in the phagosomal pathway would not undergo rigorous competition with the large pool of endogenous peptides for association with newly assembled class I MHC complexes, as would occur in the ER. It has indeed been shown that all relevant components of class I MHC loading complexes are present in early phagosomes and that these are functional [47]. How the necessary components are transported from the ER to the phagosomes is not clear. Phagosome–ER fusion was proposed [67], but other groups were unable to confirm these findings [64,68]. The vacuolar pathway is an alternative model that is based on notions of proteasome- and TAP transporter-independent cross-presentation, enabled by proteases that reside in late endosomes and lysosomes [49,69–71]. Most antigen cross-presentation studies performed in human DCs to date focus on this pathway, and less on proteasome/TAP-dependent mechanisms [58,59,71]. As only peptide-bound class I MHC molecules are transported to the plasma membrane, peptide-exchange should be able to occur in the endosomal encountered class I MHC molecules. Earlier *in vitro* experiments have already suggested that peptide-receptive class I MHC molecules can be generated under late endosomal/lysosomal pH conditions [72]. Multiple pathways can co-exist in the same cell type, indicating that these pathways are compartmentalized and require sorting and specific antigen targeting to specialized endosomal compartments [49].

Antigen uptake routes by DCs control antigen cross-presentation efficiency

To allow for the induction of specific adaptive immunity, pathogens or antigenic components that are pathogen-derived must be internalized by DCs for antigen processing and display as peptide/MHC complexes at the DC surface. It is clear that DCs can (cross-)present exogenous antigen without being infected [13]. Antigen can be acquired directly from the surrounding milieu, or can be received by a cross-

Table 1. Receptors involved in targeting antigen to the class I or class II major histocompatibility complex (MHC) antigen presentation pathway in mouse and human dendritic cells [26,38,39,41,75].

	Mouse	Human
Activating Fc receptors		
Fc γ receptors (Fc γ R)		
Fc γ I (CD64)	} MHC I [57,76–80]	} MHC I [81]
Fc γ II (CD32)		
Fc γ III (CD16)		
Pathogen recognition receptors (PRRs)		
C-type lectin receptors (CLRs)		
Type I		
Mannose receptor (MR/CD206)	MHC I [45]	MHC II [82]
DEC205 (CD205)	MHC I [23,80]	MHC I [83,84]
Type II		
DC-SIGN (CD209)	n.a.	MHC I [85,86]
Langerin (CD207)	MHC II/ MHC I [80]	n.d.
DCIR (CLEC 4A)	MHC II [46,80]	MHC I [87]
Dectin I	MHC II [88]	MHC I (CLEC 9A) [27]
Dectin II	MHC I [89]	n.d.
LOX-1	MHC I [90]	n.d.
Scavenger receptors (SR)		
SR-A1 and SR-A2	n.d.	n.d.
SR-B1 (CD36)	n.d.	MHC I [91]

The involvement of complement-receptors and Toll-like receptors has not been determined. n.d., not determined; n.a., not applicable.

presenting DC from a distant site through transport by migratory DCs. It has been shown that skin-derived migratory DCs transfer antigen to lymph node-resident DCs for efficient cross-presentation [73]. Secondly, it was shown that tumours secrete exosomes that contain proteins, which can be taken up by DCs. This system can facilitate anti-tumour immunity [12]. Thirdly, DCs use gap junctions to gain peptide antigens from adjacent cells. These peptides can be used thereafter for cross-presentation [74].

To allow for antigen internalization, DCs are equipped with a variety of receptors that can either directly recognize pathogen-associated molecular patterns (PAMPS) or indirectly via plasma complement (activated large proteolytic fragments of complement proteins, C3b, C4b, iC3b and C3d – collectively called C') that binds to complement receptors (CR1/CD35 and CR2/CD21). Immunoglobulins (Ig) present in plasma bind the immunoglobulin receptors (activating receptors FcRI, IIA and III and the inhibitory FcRIIB). Both complement fragments and Ig are soluble receptors present in plasma that bind structures on pathogen surfaces to facilitate pathogen opsonization, internalization and destruction. Secondly, small proteolytic complement protein fragments (i.e. C3a, C4a, C5a) act as chemoattractants to recruit and activate new phagocytes. CRs and FcRs allow for internalization after binding C' - or Ig-opsonized antigens. Antigen opsonization with Ig rather than C' facilitates antigen cross-presentation [57], thus supporting a role for Ig rather than C' in tailoring appropriate antigen-specific adaptive immune responses.

DCs use multiple additional membrane-expressed receptors for the internalization of antigens. The presence and

dominance of these receptors differs between DC subpopulations within species and between DC subpopulations, as compared between mouse and human [26]. Targeting specific receptors can drive the immune response either towards class II MHC-restricted CD4⁺ T helper cell responses or to class I MHC-restricted CD8⁺ cytotoxic T cell responses via cross-presentation, and can therefore be an effective method for inducing anti-viral or anti-tumour CTL responses [46]. In both mice and humans, the presence of many different uptake receptors has been shown (Table 1).

Uptake via distinct endocytic receptors controls the efficiency of cross-presentation of peptide/class I MHC complexes to CD8⁺ T cells. The effects of individual uptake receptors on antigen targeting to the class I or class II MHC presentation route seem to be roughly conserved between mice and humans, but opposing effects of some receptors related to endosomal targeting and processing of antigens have been found. In mice, antigen cross-presentation is promoted when antigen uptake occurs via MR [45], DEC205 [80,83,88,92,93], dectin-2 [89], DNGR-1 [94], Fc γ R [57,76–80] and LOX-1 [90]. Also in human cells, antigen cross-presentation is promoted upon antigen uptake via DEC-205 [83,84], DC-SIGN [85,86] and Fc γ R [81] *in vitro* as well as *in vivo* MR targeting in a humanized mouse model [95]. However, in contrast to the results in mice, MR-mediated antigen uptake induced CD4⁺ T cell responses by human DCs [82]. Antigen targeting to dectin-1 [88], DCIR-2 [46,80] and CD40 [96] induce CD4⁺ T cell responses in the mouse. Langerin (CD207)-targeted uptake induces both CD4⁺ and CD8⁺ T cell responses [80]. In humans, antigen cross-presentation is favoured by uptake via DCIR

[87]. These examples underscore that knowledge on receptor-mediated cross-presentation in mice cannot always be translated immediately to the human system. Because enhancing cross-presentation can be an effective means to improve CTL responses in diverse DC vaccination programmes, more research about receptor targeting in the human system is needed.

Antigen uptake routes in DCs can be decisive in induction of immunity or tolerance

Cytotoxic CD8⁺ T cells directed to virus-infected cells are considered crucial for efficient anti-viral responses. In parallel, the elicitation of tumour-directed CTLs is considered crucial for effective anti-tumour responses to occur. Considering tumour-associated antigens, dead tumour cells are a major antigen source for APCs [91]. For example, Asano *et al.* showed that dead tumour cells traffic via the lymph vessels to the tumour-draining lymph node where dead tumour cell-associated antigens are internalized by APCs and cross-presented to CD8⁺ T cells [97]. In cancer therapy, many investigators have taken advantage of the immunogenicity of tumour-associated antigens for tumour vaccination, either by direct injection of dead tumour cells [98] or using DCs loaded with dead tumour cells [99].

For pathogen-associated antigens, a large pool of antigens is also available in cells that are dead or dying as a consequence of the pathogen infection, forming a rich source of antigens for loading into the cross-presentation pathway. It is demonstrated that cells dying from infection are engulfed by APCs for CD8⁺ T cell activation by cross-presentation. For example, virally infected dying cells such as influenza A [91], Epstein–Barr virus [100] and canarypox virus [101] or bacterially infected dying cells from *Salmonella typhimurium* [102] induce CD8⁺ T cell responses. It is clear from these examples that providing antigen in the form of dead cells can be a powerful tool to favour cross-presentation of the antigen.

However, not all dead cells are immunogenic and induce cross-presentation. In humans, it is estimated that under homeostatic conditions approximately 1 million cells turn over each second, which does not generally result in autoreactivity [103]. However, deficiencies in the clearance of these dead cells can result in autoimmune disorders (i.e. systemic lupus erythematosus in individuals lacking early components of the complement cascades), indicating its role in maintenance of self-tolerance. Clearly, this example shows that the immune system is able to process dead cells in a tolerogenic or immunogenic manner, depending on several factors. As reviewed by Green *et al.*, these factors are related to the type of cell death, the cell death pathway, how the dead cells are engulfed, the engulfing cell, where the engulfment takes place and which cells of the immune system eventually encounter the antigens presented along with the dead cells [104]. Clarification (and possibly modulation) of these pro-

cesses should provide a venue for development of efficient cross-presentation routes that can be exploited in DC vaccination strategies. Several such attempts are exemplified in experiments on heat shock protein (HSP)-associated antigen uptake. HSPs are intracellular chaperone molecules that associate readily with neighbouring proteins, such as with antigen inside tumour cells. Injection with HSP, e.g. HSP 70, HSP 90 and glycoprotein (gp)96 induces CTL responses against the cells from which the HSPs were isolated (reviewed in [105]). While HSPs may not be essential for antigen cross-presentation, they have been shown to promote antigen cross-presentation using *in vitro* assays employing multiple cell lines as well as primary mouse and human immune cells [105–109]. The myeloid differentiation factor 88 (MyD88) is essential in the developmental maturation of DCs that allows them to prime CD8⁺ T cells through cross-presentation after uptake of HSP-coupled antigen [63].

Clinical experience with anti-tumour and anti-viral DC-vaccines

In recent years, multiple insights were obtained in mechanisms that underlie cross-presentation in mouse as well as human cells. The primary cross-presenting CD8 α ⁺ DC in mice was identified, and groups around the world are currently investigating what seems to be a human homologue, the human CD141⁺ DC. Distinct pathways are shown to be present in the murine system, and more knowledge is being increasingly gathered about the cross-presentation pathways in the human system. These mechanisms are not always conserved between species, which alerts us that knowledge gathered in mouse systems cannot be extrapolated to the human system without risk.

While our knowledge in antigen presentation biology increases, the potential benefit of its function is being explored in patients. In the last 15 years, at least 50 Phase I and Phase II trials in humans were performed using DC vaccination as anti-tumour [45,110–118] or anti-viral treatment [119,120] related to allogeneic stem cell transplantation (SCT) in both adult and paediatric settings (references [113,115–118] and [110–112,114,120], respectively). The main focus of these Phase I/II studies was safety, and none reported serious direct side effects. Although the studies were not set up for evaluation of effectiveness of induction of CTL responses, disease regression/prevention and immune responses were measured. In general, anti-tumour responses were minimal [110,112,114,117], as reviewed by Rosenberg *et al.* [121]. One recent study performed in 10 AML patients after at least one anti-leukaemic chemotherapeutic regimen, but not end-stage disease, showed more promising results. In this study, vaccination with DCs loaded with mRNA encoding Wilms' tumour 1 (WT1) protein induced complete remission in three of 10 patients and temporary remission in two additional patients. Moreover, an increased frequency of

WT1-specific CD8⁺ T cells was found in two of five tested HLA-A0201⁺ patients, correlating significantly with long-term response [118].

There are more examples of DC vaccination studies in which the induction of T cell responses was measured. For example, in studies in stage IV melanoma patients, peptide-pulsed CD34⁺ progenitor cell-derived DCs induced both CD4⁺ and CD8⁺ T cell responses [115,116]. Also, in patients suffering from breast cancer, DCs loaded exogenously with peptides derived from the human epidermal growth factor 2 (HER-2)/neu proto-oncogene and the epithelial mucin MUC1 lead to an induction of peptide-specific CTLs and decrease in serum amount of tumour markers [113]. DC vaccination in 35 non-Hodgkin B cell lymphoma patients targeting tumour-specific immunoglobulin resulted in a tumour regression rate of 31.6% [122]. In studies aimed at the induction of anti-viral immunity after allogeneic SCT, targeted mainly at human cytomegalovirus (HCMV), vaccination with pp65-pulsed DCs induced a sustained antigen-specific CD4⁺ T cell response; CTL responses were not assessed [120]. HCMV peptide-loaded DCs induced HCMV-specific CTL responses in five of 24 SCT patients at risk for HCMV after allogeneic SCT [119].

As described, the results from anti-tumour and anti-viral clinical trials show a modest immunological response, which may not yet result in an increase in patient survival. As most of these studies were designed as Phase I and Phase II safety studies the included patient groups tend to be late in disease progression (with a relatively high residual tumour load or already with virus-associated disease). Patients with a substantially lower tumour load and absent viral disease may be more likely to show benefit from induced specific anti-tumour or anti-viral activity, as also suggested by recent results [118]. Conversely, the potency of immunological responses was far from optimal, and provides opportunities for improvement. New information on human DC subsets and education of DCs allow for the optimization and improvement of current DC vaccination strategies. Distinct DC subsets offer unique possibilities in DC vaccination strategies [123]. In addition, priming DCs in a specific way determines the immunological outcome, which can be accomplished by inclusion in vaccine formulations of ligands to TLRs [124] and non-obese diabetic-like receptors (NLRs) [125] and may work through modulation of the DCs migratory and T cell stimulatory capacity. Secondly, in reported studies thus far, the elicitation or reactivation of CD4⁺ T cell responses can often be observed, and even humoral anti-tumour responses can be measured. Class I MHC-restricted CD8⁺ T cell responses are the focus in only a minority of papers, while it is CTLs that are key players in anti-tumour and anti-viral immunity. Therefore, the induction of potent antigen cross-presentation should be especially explored in current human immunology studies. Clarification of the mechanisms that increase the amount of peptide/class I MHC complexes is pivotal for the develop-

ment of next-generation DC-based anti-tumour and anti-viral intervention strategies.

Acknowledgements

We acknowledge the Wilhelmina Research Fund for financial support (to M. B.). We thank Michiel de Jong (<http://www.TomatoesandPotatoes.com>) for help with design Fig. 1. We additionally thank Sytze de Roock and Annemieke Laarhoven for critical reading of the manuscript, and the members of the Boes Laboratory and Center for Molecular and Cellular Intervention (CMCI) for helpful discussions.

Disclosure

The authors declare no conflicts of interest.

References

- 1 Banchereau J, Steinman RM. Dendritic cells and the control of immunity. *Nature* 1998; **392**:245–52.
- 2 Steinman RM. Dendritic cells and the control of immunity: enhancing the efficiency of antigen presentation. *Mt Sinai J Med* 2001; **68**:160–6.
- 3 Steinman RM, Banchereau J. Taking dendritic cells into medicine. *Nature* 2007; **449**:419–26.
- 4 Bevan MJ. Cross-priming for a secondary cytotoxic response to minor H antigens with H-2 congenic cells which do not cross-react in the cytotoxic assay. *J Exp Med* 1976; **143**:1283–8.
- 5 Heath WR, Carbone FR. Cross-presentation in viral immunity and self-tolerance. *Nat Rev Immunol* 2001; **1**:126–34.
- 6 Kurts C, Robinson BW, Knolle PA. Cross-priming in health and disease. *Nat Rev Immunol* 2010; **10**:403–14.
- 7 Sigal LJ, Crotty S, Andino R, Rock KL. Cytotoxic T-cell immunity to virus-infected non-haematopoietic cells requires presentation of exogenous antigen. *Nature* 1999; **398**:77–80.
- 8 Merckenslager M, Power MO, Pircher H, Fisher AG. Intrathymic deletion of MHC class I-restricted cytotoxic T cell precursors by constitutive cross-presentation of exogenous antigen. *Eur J Immunol* 1999; **29**:1477–86.
- 9 Kurts C, Kosaka H, Carbone FR, Miller JF, Heath WR. Class I-restricted cross-presentation of exogenous self-antigens leads to deletion of autoreactive CD8(+) T cells. *J Exp Med* 1997; **186**:239–45.
- 10 Luckashenak N, Schroeder S, Endt K *et al.* Constitutive cross-presentation of tissue antigens by dendritic cells controls CD8⁺ T cell tolerance *in vivo*. *Immunity* 2008; **28**:521–32.
- 11 Kurts C, Heath WR, Carbone FR, Allison J, Miller JF, Kosaka H. Constitutive class I-restricted exogenous presentation of self antigens *in vivo*. *J Exp Med* 1996; **184**:923–30.
- 12 Wolfers J, Lozier A, Raposo G *et al.* Tumor-derived exosomes are a source of shared tumor rejection antigens for CTL cross-priming. *Nat Med* 2001; **7**:297–303.
- 13 Albert ML, Sauter B, Bhardwaj N. Dendritic cells acquire antigen from apoptotic cells and induce class I-restricted CTLs. *Nature* 1998; **392**:86–9.
- 14 Kovacsics-Bankowski M, Clark K, Benacerraf B, Rock KL. Effi-

- cient major histocompatibility complex class I presentation of exogenous antigen upon phagocytosis by macrophages. *Proc Natl Acad Sci USA* 1993; **90**:4942–6.
- 15 Shen Z, Reznikoff G, Dranoff G, Rock KL. Cloned dendritic cells can present exogenous antigens on both MHC class I and class II molecules. *J Immunol* 1997; **158**:2723–30.
 - 16 Limmer A, Ohl J, Kurts C *et al*. Efficient presentation of exogenous antigen by liver endothelial cells to CD8+ T cells results in antigen-specific T-cell tolerance. *Nat Med* 2000; **6**:1348–54.
 - 17 Heit A, Huster KM, Schmitz F, Schiemann M, Busch DH, Wagner H. CpG-DNA aided cross-priming by cross-presenting B cells. *J Immunol* 2004; **172**:1501–7.
 - 18 Hon H, Oran A, Brocker T, Jacob J. B lymphocytes participate in cross-presentation of antigen following gene gun vaccination. *J Immunol* 2005; **174**:5233–42.
 - 19 Tvinnereim AR, Hamilton SE, Harty JT. Neutrophil involvement in cross-priming CD8+ T cell responses to bacterial antigens. *J Immunol* 2004; **173**:1994–2002.
 - 20 Beauvillain C, Delneste Y, Scotet M *et al*. Neutrophils efficiently cross-prime naive T cells *in vivo*. *Blood* 2007; **110**:2965–73.
 - 21 Kim S, Shen T, Min B. Basophils can directly present or cross-present antigen to CD8 lymphocytes and alter CD8 T cell differentiation into IL-10-producing phenotypes. *J Immunol* 2009; **183**:3033–9.
 - 22 Stelekati E, Bahri R, D'Orlando O *et al*. Mast cell-mediated antigen presentation regulates CD8+ T cell effector functions. *Immunity* 2009; **31**:665–76.
 - 23 Bagai R, Valujskikh A, Canaday DH *et al*. Mouse endothelial cells cross-present lymphocyte-derived antigen on class I MHC via a TAP1- and proteasome-dependent pathway. *J Immunol* 2005; **174**:7711–15.
 - 24 Jung S, Unutmaz D, Wong P *et al*. *In vivo* depletion of CD11c+ dendritic cells abrogates priming of CD8+ T cells by exogenous cell-associated antigens. *Immunity* 2002; **17**:211–20.
 - 25 Storni T, Bachmann MF. Loading of MHC class I and II presentation pathways by exogenous antigens: a quantitative *in vivo* comparison. *J Immunol* 2004; **172**:6129–35.
 - 26 Hart DN. Dendritic cells: unique leukocyte populations which control the primary immune response. *Blood* 1997; **90**:3245–87.
 - 27 Jongbloed SL, Kassianos AJ, McDonald KJ *et al*. Human CD141+ (BDCA-3)+ dendritic cells (DCs) represent a unique myeloid DC subset that cross-presents necrotic cell antigens. *J Exp Med* 2010; **207**:1247–60.
 - 28 MacDonald KP, Munster DJ, Clark GJ, Dzionek A, Schmitz J, Hart DN. Characterization of human blood dendritic cell subsets. *Blood* 2002; **100**:4512–20.
 - 29 Vremec D, Zorbas M, Scollay R *et al*. The surface phenotype of dendritic cells purified from mouse thymus and spleen: investigation of the CD8 expression by a subpopulation of dendritic cells. *J Exp Med* 1992; **176**:47–58.
 - 30 del Rio ML, Rodriguez-Barbosa JI, Kremmer E, Forster R. CD103+ and CD103+ bronchial lymph node dendritic cells are specialized in presenting and cross-presenting innocuous antigen to CD4+ and CD8+ T cells. *J Immunol* 2007; **178**:6861–6.
 - 31 den Haan JM, Lehar SM, Bevan MJ. CD8(+) but not CD8(-) dendritic cells cross-prime cytotoxic T cells *in vivo*. *J Exp Med* 2000; **192**:1685–96.
 - 32 Iyoda T, Shimoyama S, Liu K *et al*. The CD8+ dendritic cell subset selectively endocytoses dying cells in culture and *in vivo*. *J Exp Med* 2002; **195**:1289–302.
 - 33 Naik SH, Proietto AI, Wilson NS *et al*. Cutting edge: generation of splenic CD8+ and CD8- dendritic cell equivalents in Fms-like tyrosine kinase 3 ligand bone marrow cultures. *J Immunol* 2005; **174**:6592–7.
 - 34 Schnorrer P, Behrens GM, Wilson NS *et al*. The dominant role of CD8+ dendritic cells in cross-presentation is not dictated by antigen capture. *Proc Natl Acad Sci USA* 2006; **103**:10729–34.
 - 35 Schulz O, Reis e Sousa C. Cross-presentation of cell-associated antigens by CD8alpha+ dendritic cells is attributable to their ability to internalize dead cells. *Immunology* 2002; **107**:183–9.
 - 36 Valdez Y, Mah W, Winslow MM, Xu L, Ling P, Townsend SE. Major histocompatibility complex class II presentation of cell-associated antigen is mediated by CD8alpha+ dendritic cells *in vivo*. *J Exp Med* 2002; **195**:683–94.
 - 37 Hildner K, Edelson BT, Purtha WE *et al*. Batf3 deficiency reveals a critical role for CD8alpha+ dendritic cells in cytotoxic T cell immunity. *Science* 2008; **322**:1097–100.
 - 38 Caminschi I, Proietto AI, Ahmet F *et al*. The dendritic cell subtype-restricted C-type lectin Clec9A is a target for vaccine enhancement. *Blood* 2008; **112**:3264–73.
 - 39 Huysamen C, Willment JA, Dennehy KM, Brown GD. CLEC9A is a novel activation C-type lectin-like receptor expressed on BDCA3+ dendritic cells and a subset of monocytes. *J Biol Chem* 2008; **283**:16693–701.
 - 40 Bachem A, Guttler S, Hartung E *et al*. Superior antigen cross-presentation and XCR1 expression define human CD11c+CD141+ cells as homologues of mouse CD8+ dendritic cells. *J Exp Med* 2010; **207**:1273–81.
 - 41 Poulin LF, Salio M, Griessinger E *et al*. Characterization of human DNGR-1+ BDCA3+ leukocytes as putative equivalents of mouse CD8alpha+ dendritic cells. *J Exp Med* 2010; **207**:1261–71.
 - 42 Savina A, Peres A, Cebrian I *et al*. The small GTPase Rac2 controls phagosomal alkalinization and antigen crosspresentation selectively in CD8(+) dendritic cells. *Immunity* 2009; **30**:544–55.
 - 43 Delamarre L, Pack M, Chang H, Mellman I, Trombetta ES. Differential lysosomal proteolysis in antigen-presenting cells determines antigen fate. *Science* 2005; **307**:1630–4.
 - 44 Trombetta ES, Ebersold M, Garrett W, Pypaert M, Mellman I. Activation of lysosomal function during dendritic cell maturation. *Science* 2003; **299**:1400–3.
 - 45 Burgdorf S, Lukacs-Kornek V, Kurts C. The mannose receptor mediates uptake of soluble but not of cell-associated antigen for cross-presentation. *J Immunol* 2006; **176**:6770–6.
 - 46 Dudziak D, Kamphorst AO, Heidkamp GF *et al*. Differential antigen processing by dendritic cell subsets *in vivo*. *Science* 2007; **315**:107–11.
 - 47 Ackerman AL, Kyritsis C, Tampe R, Cresswell P. Access of soluble antigens to the endoplasmic reticulum can explain cross-presentation by dendritic cells. *Nat Immunol* 2005; **6**:107–13.
 - 48 Kovacsics-Bankowski M, Rock KL. A phagosome-to-cytosol pathway for exogenous antigens presented on MHC class I molecules. *Science* 1995; **267**:243–6.
 - 49 Shen L, Sigal LJ, Boes M, Rock KL. Important role of cathepsin S in generating peptides for TAP-independent MHC class I cross-presentation *in vivo*. *Immunity* 2004; **21**:155–65.
 - 50 Song R, Harding CV. Roles of proteasomes, transporter for antigen presentation (TAP), and beta 2-microglobulin in the processing of bacterial or particulate antigens via an alternate class I MHC processing pathway. *J Immunol* 1996; **156**:4182–90.
 - 51 Lennon-Dumenil AM, Bakker AH, Maehr R *et al*. Analysis of pro-

- tease activity in live antigen-presenting cells shows regulation of the phagosomal proteolytic contents during dendritic cell activation. *J Exp Med* 2002; **196**:529–40.
- 52 Kinchen JM, Ravichandran KS. Phagosome maturation: going through the acid test. *Nat Rev Mol Cell Biol* 2008; **9**:781–95.
- 53 Kawane K, Ohtani M, Miwa K *et al.* Chronic polyarthritis caused by mammalian DNA that escapes from degradation in macrophages. *Nature* 2006; **443**:998–1002.
- 54 Savina A, Jancic C, Hugues S *et al.* NOX2 controls phagosomal pH to regulate antigen processing during crosspresentation by dendritic cells. *Cell* 2006; **126**:205–18.
- 55 Yates RM, Russell DG. Phagosome maturation proceeds independently of stimulation of toll-like receptors 2 and 4. *Immunity* 2005; **23**:409–17.
- 56 Jancic C, Savina A, Wasmeier C *et al.* Rab27a regulates phagosomal pH and NADPH oxidase recruitment to dendritic cell phagosomes. *Nat Cell Biol* 2007; **9**:367–78.
- 57 Kim SH, Visser A, Crujisen C, van d V, Boes M. Recruitment of Rab27a to phagosomes controls microbial antigen cross-presentation by dendritic cells. *Infect Immun* 2008; **76**:5373–80.
- 58 Saveanu L, Carroll O, Weimershaus M *et al.* IRAP identifies an endosomal compartment required for MHC class I cross-presentation. *Science* 2009; **325**:213–17.
- 59 Segura E, Albiston AL, Wicks IP, Chai SY, Villadangos JA. Different cross-presentation pathways in steady-state and inflammatory dendritic cells. *Proc Natl Acad Sci USA* 2009; **106**:20377–81.
- 60 Ackerman AL, Giodini A, Cresswell P. A role for the endoplasmic reticulum protein retrotranslocation machinery during crosspresentation by dendritic cells. *Immunity* 2006; **25**:607–17.
- 61 Rodriguez A, Regnault A, Kleijmeer M, Ricciardi-Castagnoli P, Amigorena S. Selective transport of internalized antigens to the cytosol for MHC class I presentation in dendritic cells. *Nat Cell Biol* 1999; **1**:362–8.
- 62 Singh R, Cresswell P. Defective cross-presentation of viral antigens in GILT-free mice. *Science* 2010; **328**:1394–8.
- 63 Palliser D, Ploegh H, Boes M. Myeloid differentiation factor 88 is required for cross-priming *in vivo*. *J Immunol* 2004; **172**:3415–21.
- 64 Burgdorf S, Scholz C, Kautz A, Tampe R, Kurts C. Spatial and mechanistic separation of cross-presentation and endogenous antigen presentation. *Nat Immunol* 2008; **9**:558–66.
- 65 Ackerman AL, Kyritsis C, Tampe R, Cresswell P. Early phagosomes in dendritic cells form a cellular compartment sufficient for cross presentation of exogenous antigens. *Proc Natl Acad Sci USA* 2003; **100**:12889–94.
- 66 Houde M, Bertholet S, Gagnon E *et al.* Phagosomes are competent organelles for antigen cross-presentation. *Nature* 2003; **425**:402–6.
- 67 Guernonprez P, Saveanu L, Kleijmeer M, Davoust J, van EP, Amigorena S. ER-phagosome fusion defines an MHC class I cross-presentation compartment in dendritic cells. *Nature* 2003; **425**:397–402.
- 68 Touret N, Paroutis P, Terebiznik M *et al.* Quantitative and dynamic assessment of the contribution of the ER to phagosome formation. *Cell* 2005; **123**:157–70.
- 69 Campbell DJ, Serwold T, Shastri N. Bacterial proteins can be processed by macrophages in a transporter associated with antigen processing-independent, cysteine protease-dependent manner for presentation by MHC class I molecules. *J Immunol* 2000; **164**:168–75.
- 70 Pfeifer JD, Wick MJ, Roberts RL, Findlay K, Normark SJ, Harding CV. Phagocytic processing of bacterial antigens for class I MHC presentation to T cells. *Nature* 1993; **361**:359–62.
- 71 Di Pucchio T, Chatterjee B, Smed-Sorensen A *et al.* Direct proteasome-independent cross-presentation of viral antigen by plasmacytoid dendritic cells on major histocompatibility complex class I. *Nat Immunol* 2008; **9**:551–7.
- 72 Gromme M, Uytdehaag FG, Janssen H *et al.* Recycling MHC class I molecules and endosomal peptide loading. *Proc Natl Acad Sci USA* 1999; **96**:10326–31.
- 73 Allan RS, Waithman J, Bedoui S *et al.* Migratory dendritic cells transfer antigen to a lymph node-resident dendritic cell population for efficient CTL priming. *Immunity* 2006; **25**:153–62.
- 74 Neijssen J, Pang B, Neefjes J. Gap junction-mediated intercellular communication in the immune system. *Prog Biophys Mol Biol* 2007; **94**:207–18.
- 75 Figdor CG, van KY, Adema GJ. C-type lectin receptors on dendritic cells and Langerhans cells. *Nat Rev Immunol* 2002; **2**:77–84.
- 76 Amigorena S, Bonnerot C. Fc receptors for IgG and antigen presentation on MHC class I and class II molecules. *Semin Immunol* 1999; **11**:385–90.
- 77 Amigorena S. Fc gamma receptors and cross-presentation in dendritic cells. *J Exp Med* 2002; **195**:F1–F3.
- 78 den Haan JM, Bevan MJ. Constitutive versus activation-dependent cross-presentation of immune complexes by CD8(+) and CD8(-) dendritic cells *in vivo*. *J Exp Med* 2002; **196**:817–27.
- 79 Kalergis AM, Ravetch JV. Inducing tumor immunity through the selective engagement of activating Fc gamma receptors on dendritic cells. *J Exp Med* 2002; **195**:1653–9.
- 80 Regnault A, Lankar D, Lacabanne V *et al.* Fc gamma receptor-mediated induction of dendritic cell maturation and major histocompatibility complex class I-restricted antigen presentation after immune complex internalization. *J Exp Med* 1999; **189**:371–80.
- 81 Liu Y, Gao X, Masuda E, Redecha PB, Blank MC, Pricop L. Regulated expression of Fc gammaR in human dendritic cells controls cross-presentation of antigen-antibody complexes. *J Immunol* 2006; **177**:8440–7.
- 82 Tan MC, Mommaas AM, Drijfhout JW *et al.* Mannose receptor-mediated uptake of antigens strongly enhances HLA class II-restricted antigen presentation by cultured dendritic cells. *Eur J Immunol* 1997; **27**:2426–35.
- 83 Bozzacco L, Trumpfheller C, Siegal FP *et al.* DEC-205 receptor on dendritic cells mediates presentation of HIV gag protein to CD8+ T cells in a spectrum of human MHC I haplotypes. *Proc Natl Acad Sci USA* 2007; **104**:1289–94.
- 84 Tsuji T, Matsuzaki J, Kelly MP *et al.* Antibody-targeted NY-ESO-1 to mannose receptor or DEC-205 *in vitro* elicits dual human CD8+ and CD4+ T cell responses with broad antigen specificity. *J Immunol* 2010; **186**:1218–27.
- 85 Tacke PJ, de V I, Gijzen K *et al.* Effective induction of naive and recall T-cell responses by targeting antigen to human dendritic cells via a humanized anti-DC-SIGN antibody. *Blood* 2005; **106**:1278–85.
- 86 Tacke PJ, Joosten B, Reddy A *et al.* No advantage of cell-penetrating peptides over receptor-specific antibodies in targeting antigen to human dendritic cells for cross-presentation. *J Immunol* 2008; **180**:7687–96.
- 87 Klechevsky E, Flamar AL, Cao Y *et al.* Cross-priming CD8+ T cells by targeting antigens to human dendritic cells through DCIR. *Blood* 2010; **116**:1685–97.

- 88 Carter RW, Thompson C, Reid DM, Wong SY, Tough DF. Preferential induction of CD4+ T cell responses through *in vivo* targeting of antigen to dendritic cell-associated C-type lectin-1. *J Immunol* 2006; **177**:2276–84.
- 89 Carter RW, Thompson C, Reid DM, Wong SY, Tough DF. Induction of CD8+ T cell responses through targeting of antigen to Dectin-2. *Cell Immunol* 2006; **239**:87–91.
- 90 Delneste Y, Magistrelli G, Gauchat J *et al.* Involvement of LOX-1 in dendritic cell-mediated antigen cross-presentation. *Immunity* 2002; **17**:353–62.
- 91 Albert ML, Pearce SF, Francisco LM *et al.* Immature dendritic cells phagocytose apoptotic cells via α v β 5 and CD36, and cross-present antigens to cytotoxic T lymphocytes. *J Exp Med* 1998; **188**:1359–68.
- 92 Bozzacco L, Trumpfheller C, Huang Y *et al.* HIV gag protein is efficiently cross-presented when targeted with an antibody towards the DEC-205 receptor in Flt3 ligand-mobilized murine DC. *Eur J Immunol* 2010; **40**:36–46.
- 93 Bonifaz L, Bonnyay D, Mahnke K, Rivera M, Nussenzweig MC, Steinman RM. Efficient targeting of protein antigen to the dendritic cell receptor DEC-205 in the steady state leads to antigen presentation on major histocompatibility complex class I products and peripheral CD8+ T cell tolerance. *J Exp Med* 2002; **196**:1627–38.
- 94 Sancho D, Mourao-Sa D, Joffre OP *et al.* Tumor therapy in mice via antigen targeting to a novel, DC-restricted C-type lectin. *J Clin Invest* 2008; **118**:2098–110.
- 95 He LZ, Crocker A, Lee J *et al.* Antigenic targeting of the human mannose receptor induces tumor immunity. *J Immunol* 2007; **178**:6259–67.
- 96 Schjetne KW, Fredriksen AB, Bogen B. Delivery of antigen to CD40 induces protective immune responses against tumors. *J Immunol* 2007; **178**:4169–76.
- 97 Asano K, Nabeyama A, Miyake Y *et al.* CD169-positive macrophages dominate antitumor immunity by crosspresenting dead cell-associated antigens. *Immunity* 2011; **34**:85–95.
- 98 Obeid M, Tesniere A, Ghiringhelli F *et al.* Calreticulin exposure dictates the immunogenicity of cancer cell death. *Nat Med* 2007; **13**:54–61.
- 99 Berard F, Blanco P, Davoust J *et al.* Cross-priming of naive CD8 T cells against melanoma antigens using dendritic cells loaded with killed allogeneic melanoma cells. *J Exp Med* 2000; **192**:1535–44.
- 100 Subklewe M, Paludan C, Tsang ML, Mahnke K, Steinman RM, Munz C. Dendritic cells cross-present latency gene products from Epstein–Barr virus-transformed B cells and expand tumor-reactive CD8(+) killer T cells. *J Exp Med* 2001; **193**:405–11.
- 101 Motta I, Andre F, Lim A *et al.* Cross-presentation by dendritic cells of tumor antigen expressed in apoptotic recombinant canarypox virus-infected dendritic cells. *J Immunol* 2001; **167**:1795–802.
- 102 Yrlid U, Wick MJ. Salmonella-induced apoptosis of infected macrophages results in presentation of a bacteria-encoded antigen after uptake by bystander dendritic cells. *J Exp Med* 2000; **191**:613–24.
- 103 Reed JC. Drug insight: cancer therapy strategies based on restoration of endogenous cell death mechanisms. *Nat Clin Pract Oncol* 2006; **3**:388–98.
- 104 Green DR, Ferguson T, Zitvogel L, Kroemer G. Immunogenic and tolerogenic cell death. *Nat Rev Immunol* 2009; **9**:353–63.
- 105 Li Z, Menoret A, Srivastava P. Roles of heat-shock proteins in antigen presentation and cross-presentation. *Curr Opin Immunol* 2002; **14**:45–51.
- 106 Ichiiyanagi T, Imai T, Kajiwaru C *et al.* Essential role of endogenous heat shock protein 90 of dendritic cells in antigen cross-presentation. *J Immunol* 2010; **185**:2693–700.
- 107 Castellino F, Boucher PE, Eichelberg K *et al.* Receptor-mediated uptake of antigen/heat shock protein complexes results in major histocompatibility complex class I antigen presentation via two distinct processing pathways. *J Exp Med* 2000; **191**:1957–64.
- 108 Singh-Jasuja H, Toes RE, Spee P *et al.* Cross-presentation of glycoprotein 96-associated antigens on major histocompatibility complex class I molecules requires receptor-mediated endocytosis. *J Exp Med* 2000; **191**:1965–74.
- 109 Callahan MK, Garg M, Srivastava PK. Heat-shock protein 90 associates with N-terminal extended peptides and is required for direct and indirect antigen presentation. *Proc Natl Acad Sci USA* 2008; **105**:1662–7.
- 110 Dohnal AM, Witt V, Hugel H, Holter W, Gadner H, Felzmann T. Phase I study of tumor Ag-loaded IL-12 secreting semi-mature DC for the treatment of pediatric cancer. *Cytotherapy* 2007; **9**:755–70.
- 111 Caruso DA, Orme LM, Neale AM *et al.* Results of a phase 1 study utilizing monocyte-derived dendritic cells pulsed with tumor RNA in children and young adults with brain cancer. *Neuro Oncol* 2004; **6**:236–46.
- 112 Caruso DA, Orme LM, Amor GM *et al.* Results of a Phase I study utilizing monocyte-derived dendritic cells pulsed with tumor RNA in children with Stage 4 neuroblastoma. *Cancer* 2005; **103**:1280–91.
- 113 Brossart P, Wirths S, Stuhler G, Reichardt VL, Kanz L, Brugger W. Induction of cytotoxic T-lymphocyte responses *in vivo* after vaccinations with peptide-pulsed dendritic cells. *Blood* 2000; **96**:3102–8.
- 114 Geiger JD, Hutchinson RJ, Hohenkirk LF *et al.* Vaccination of pediatric solid tumor patients with tumor lysate-pulsed dendritic cells can expand specific T cells and mediate tumor regression. *Cancer Res* 2001; **61**:8513–19.
- 115 Banchereau J, Palucka AK, Dhodapkar M *et al.* Immune and clinical responses in patients with metastatic melanoma to CD34(+) progenitor-derived dendritic cell vaccine. *Cancer Res* 2001; **61**:6451–8.
- 116 Paczesny S, Banchereau J, Wittkowski KM, Saracino G, Fay J, Palucka AK. Expansion of melanoma-specific cytolytic CD8+ T cell precursors in patients with metastatic melanoma vaccinated with CD34+ progenitor-derived dendritic cells. *J Exp Med* 2004; **199**:1503–11.
- 117 Banchereau J, Ueno H, Dhodapkar M *et al.* Immune and clinical outcomes in patients with stage IV melanoma vaccinated with peptide-pulsed dendritic cells derived from CD34+ progenitors and activated with type I interferon. *J Immunother* 2005; **28**:505–16.
- 118 Tendeloo V, van d V, Van DA *et al.* Induction of complete and molecular remissions in acute myeloid leukemia by Wilms' tumor 1 antigen-targeted dendritic cell vaccination. *Proc Natl Acad Sci USA* 2010; **107**:13824–9.
- 119 Grigoleit GU, Kapp M, Hebart H *et al.* Dendritic cell vaccination in allogeneic stem cell recipients: induction of human cytomegalovirus (HCMV)-specific cytotoxic T lymphocyte responses even in patients receiving a transplant from an HCMV-seronegative donor. *J Infect Dis* 2007; **196**:699–704.
- 120 Feuchtinger T, Opher K, Bicanic O *et al.* Dendritic cell vaccination in an allogeneic stem cell recipient receiving a transplant from a human cytomegalovirus (HCMV)-seronegative donor: induction of a HCMV-specific T(helper) cell response. *Cytotherapy* 2010; **12**:945–50.

- 121 Rosenberg SA, Yang JC, Restifo NP. Cancer immunotherapy: moving beyond current vaccines. *Nat Med* 2004; **10**:909–15.
- 122 Timmerman JM, Czerwinski DK, Davis TA *et al.* Idiotype-pulsed dendritic cell vaccination for B-cell lymphoma: clinical and immune responses in 35 patients. *Blood* 2002; **99**:1517–26.
- 123 Schreibelt G, Tel J, Sliepen KH *et al.* Toll-like receptor expression and function in human dendritic cell subsets: implications for dendritic cell-based anti-cancer immunotherapy. *Cancer Immunol Immunother* 2010; **59**:1573–82.
- 124 Jelinek I, Leonard JN, Price GE *et al.* TLR3-specific double-stranded RNA oligonucleotide adjuvants induce dendritic cell cross-presentation, CTL responses, and antiviral protection. *J Immunol* 2011; **186**:2422–9.
- 125 Asano J, Tada H, Onai N *et al.* Nucleotide oligomerization binding domain-like receptor signaling enhances dendritic cell-mediated cross-priming *in vivo*. *J Immunol* 2010; **184**:736–45.